

Coloring of graphs associated to zero-divisors

Hsin-Ju Wang ^{*†}

Abstract. Let G be a graph, $\chi(G)$ be the minimal number of colors which can be assigned to the vertices of G in such a way that every two adjacent vertices have different colors and $\omega(G)$ to be the least upper bound of the size of the complete subgraphs contained in G . It is well-known that $\chi(G) \geq \omega(G)$. Beck in [6] conjectured that $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R))$ if $\omega(\Gamma_0(R)) < \infty$, where $\Gamma_0(R)$ is a graph associated to a commutative ring R . In this note, we provide some sufficient conditions for a ring R to enjoy $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R))$. As a consequence, we verify Beck's conjecture for the homomorphic image of $\mathbb{Z}^n$.

1 Introduction

In this paper, R will denote a commutative ring with 1. If S is a subset of R , we denote $S - \{0\}$ by S^* . Also, we use $\mathbb{N}$ for the set of all nonnegative integers.

In [6], Beck introduced the idea of a *zero-divisor graph* of a commutative ring R with 1. He defined $\Gamma_0(R)$ to be the graph whose vertices are elements of R and in which two vertices x and y are adjacent if and only if $xy = 0$. Beck was mostly concerned with coloring $\Gamma_0(R)$. Recall that the *chromatic number* of the graph, denoted $\chi(\Gamma_0(R))$, is defined to be the minimal number of colors which can be assigned to the elements of $\Gamma_0(R)$ in such a way that every two adjacent elements have different colors. A subset C of vertex set of $\Gamma_0(R)$ is called a *clique* if any two distinct elements of C are adjacent; the number $\omega(\Gamma_0(R))$ is the least upper bound of the size of the cliques, and clearly $\chi(\Gamma_0(R)) \geq \omega(\Gamma_0(R))$. Beck conjectured that $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R))$ if $\omega(\Gamma_0(R)) < \infty$ and proved this conjecture for a rather wide class of rings,

^{*}**Key Words:** zero-divisor graph, chromatic number, clique number, complement graph

[†]2000 Mathematics Subject Classification. Primary 13A99, 05C15; Secondary 13M99

including reduced rings and principal ideal rings. However, D.D. Anderson and M. Naseer gave a counterexample to this conjecture in [5]. One of the aim of this paper is to find some sufficient conditions for rings R to enjoy $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R))$. As a consequence, we are able to verify the conjecture for a class of rings, namely, the homomorphic image of $\mathbb{Z}^n$.

Let R be a ring and let $Z(R)$ be the set of zero-divisors of R . Then there is associated a (simple) graph $\Gamma(R)$ to R with vertices $Z(R)^*$ and where two distinct vertices $x, y \in Z(R)^*$ are adjacent if and only if $xy = 0$. This graph was defined slightly differently than the graph introduced by I. Beck and was introduced by Anderson and Livingston in [4]. Recently, this concept was studied extensively in [1], [2], [3], [4], [9] and [10]. One of the aim of this paper is to prove that $\chi(\Gamma(R)) = \omega(\Gamma(R))$ if $R \cong \mathbb{Z}_{p_1^{r_1}} \times \cdots \times \mathbb{Z}_{p_k^{r_k}}$, where p_i is a prime number for every i and $\bar{G}$ is the complement graph of G .

2 Preliminary

We review some background from graph theory and fix some notations from [5] and [6] in this section.

A simple graph G is an ordered pair of disjoint sets (V, E) such that $V = V(G)$ is the vertex set of G and $E = E(G)$ is the edge set of G . Often we use G for $V(G)$. The *order* of a graph G , written by $|G|$, is the cardinality of $V(G)$. A *subgraph* of G is a graph having all of its vertices and edges in G . Let $V' \subseteq V(G)$; then $G - V'$ is the subgraph of G obtained by deleting the vertices in V' and all edges incident with them. For $v \in V$, the *degree* of v , denoted by $\deg(v)$, is the number of edges of G incident to v . A vertex subset S of a graph G is called an *independent set* if any two vertices of S are not adjacent.

A graph G is said to be a *totally disconnected* graph if the edge set $E(G) = \emptyset$. A graph in which each pair of distinct vertices is joined by an edge is called a *complete graph*. We use K_n for the complete graph with n vertices.

The most important result in [6] is the following.

Theorem 2.1 [6, Theorem 3.9] *The following are equivalent for a ring R :*

- (i) $\chi(\Gamma_0(R)) < \infty$.
- (ii) $\omega(\Gamma_0(R)) < \infty$.

(iii) The nil-radical is finite and equals a finite intersection of prime ideals.

Beck named the rings satisfying one of the three conditions in Theorem 2.1 as follows.

Definition 2.2 A ring R is called a coloring provided that $\chi(\Gamma_0(R)) < \infty$.

In case $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) < \infty$ for rings R , Anderson and Naseer made the following definition.

Definition 2.3 A ring R is called a chromatic provided that $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) < \infty$.

In the sequel, we use the symbol $A \sqcup B$ to denote the disjoint union of two sets A and B .

3 Coloring of $\Gamma_0(R)$

Let R be a commutative ring with 1 and let $Z(R)$ be the set of zero-divisors of R . Let $\Gamma_0(R)$ and $\Gamma(R)$ be the graphs associated to R defined in the introduction. Observe that, in $\Gamma_0(R)$, 0 is adjacent to every other vertices of $\Gamma_0(R)$ and the degree of every unit is 1. Thus $\Gamma(R)$ is a subgraph of $\Gamma_0(R)$ and $\Gamma_0(R) - \{0\}$ is a disjoint union of $\Gamma(R)$ and a totally disconnected graph.

The goal of this section is to study the interplay between $\chi(\Gamma_0(R))$ and $\omega(\Gamma_0(R))$. For this, we first look at some special cases.

Lemma 3.1 Let $R \cong \mathbb{Z}_{p^r}$, where p is a prime number and r be a positive integer. Then R is a chromatic ring with $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = p^{r/2}$ if r is even and $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = p^{(r-1)/2} + 1$ if r is odd.

Proof. Suppose that $r = 2t$ is even. Then $S = \{p^t n \mid n \in \mathbb{Z}\}$ is a clique subset of $\Gamma_0(R)$. Since $|S| = p^{r/2}$, we require $p^{r/2}$ colors to paint S . Observe that p^t is not adjacent to any vertex of $\Gamma_0(R) - S$ and $\Gamma_0(R) - S$ is an independent set. We can use the color of p^t to color every vertex of $\Gamma_0(R) - S$. Thus $\chi(\Gamma_0(R)) \leq p^{r/2} \leq \omega(\Gamma_0(R)) \leq \chi(\Gamma_0(R))$.

Suppose that $r = 2t + 1$ is odd. Then $S = \{p^{t+1} n \mid n \in \mathbb{Z}\} \cup \{p^t\}$ is a clique subset of $\Gamma_0(R)$. Since $|S| = p^{(r-1)/2} + 1$, we require $p^{(r-1)/2} + 1$ colors to paint S . Observe that p^t is not adjacent to any vertex of $\Gamma_0(R) - S$ and $\Gamma_0(R) - S$ is an independent set. We can use the color of p^t to color every

vertex of $\Gamma_0(R) - S$. Thus $\chi(\Gamma_0(R)) \leq p^{(r-1)/2} + 1 \leq \omega(\Gamma_0(R)) \leq \chi(\Gamma_0(R))$.
 $\square$

From the proof of Lemma 3.1, we have the following consequence.

Corollary 3.2 *Let $R \cong \mathbb{Z}_{p^r}$, where p is a prime number and r be a positive integer. Let S be a largest clique of $\Gamma_0(R)$; then $|\{a \in S \mid a^2 \neq 0\}| = 0$ if r is even and $|\{a \in S \mid a^2 \neq 0\}| = 1$ if r is odd.*

Example 3.3 *Let $R \cong \mathbb{Z}_8 \times \mathbb{Z}_{16}$; then $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = 9$.*

Proof. Let $C_1 = \{(0, 0), (0, 4), (0, 8), (0, 12), (4, 0), (4, 4), (4, 8), (4, 12), (2, 0)\}$, $C_2 = (\{1, 3, 5, 7\} \times \{0, 4, 8, 12\}) \cup ((\{2, 6\} \times \{0, 4, 8, 12\}) - \{(2, 0)\})$ and $C_3 = \mathbb{Z}_8 \times (\mathbb{Z}_{16} - \{0, 4, 8, 12\})$; then C_1 is a clique and $\Gamma_0(R) = C_1 \sqcup C_2 \sqcup C_3$. Moreover, C_2 and C_3 are independent sets. Observe that we require 9 colors to paint C_1 . Then we can use the color of $(2, 0)$ (resp. $(0, 4)$) to color C_2 (resp. C_3). In such a way, every two adjacent vertices of $\Gamma_0(R)$ have different colors. Thus, we conclude that $9 \leq \omega(\Gamma_0(R)) \leq \chi(\Gamma_0(R)) \leq 9$. $\square$

The following result stated in [6] without any proof. Here, we provide one.

Lemma 3.4 *Let R_1 and R_2 be rings and $R \cong R_1 \times R_2$. Then the following hold.*

- (i) $\chi(\Gamma_0(R)) \geq \chi(\Gamma_0(R_1)) + \chi(\Gamma_0(R_2)) - 1$.
- (ii) *If R_2 is reduced, then $\chi(\Gamma_0(R)) = \chi(\Gamma_0(R_1)) + \chi(\Gamma_0(R_2)) - 1$.*

Proof. (i) If $\chi(\Gamma_0(R_1))$ or $\chi(\Gamma_0(R_2))$ are infinite then so is $\chi(\Gamma_0(R))$. Therefore the inequality holds trivially. So we may assume that $t_i = \chi(\Gamma_0(R_i)) < \infty$ for $i = 1, 2$. Thus, we require t_1 colors to paint $\Gamma_0(R_1) \times \{0\}$ and t_2 colors to paint $\{0\} \times \Gamma_0(R_2)$. Since $(0, 0)$ is adjacent to any vertex of $\Gamma_0(R)$ and every vertex of $\Gamma_0(R_1) \times \{0\}$ is adjacent to every vertex of $\{0\} \times \Gamma_0(R_2)$, we require at least $t_1 + t_2 - 1$ colors to paint $\Gamma_0(R)$.

(ii) Suppose that R_2 is reduced. As in (i), we may assume that $t_i = \chi(\Gamma_0(R_i)) < \infty$ for $i = 1, 2$. Observe that $t_i - 1 = \chi(\Gamma_0(R_i) - \{0\})$ for $i = 1, 2$, there are independent sets $S_1, \dots, S_{t_1-1}$ of the graph $\Gamma_0(R_1)$ and independent sets $T_1, \dots, T_{t_2-1}$ of the graph $\Gamma_0(R_2)$ such that $\Gamma_0(R_1) - \{0\} = S_1 \sqcup \dots \sqcup S_{t_1-1}$ and $\Gamma_0(R_2) - \{0\} = T_1 \sqcup \dots \sqcup T_{t_2-1}$. Thus,

$$\begin{aligned} \Gamma_0(R) = & (0, 0) \sqcup (S_1 \times \{0\}) \sqcup \dots \sqcup (S_{t_1-1} \times \{0\}) \\ & \sqcup (\Gamma_0(R_1) \times T_1) \sqcup \dots \sqcup (\Gamma_0(R_1) \times T_{t_2-1}). \end{aligned} \quad (1)$$

Observe that $\Gamma_0(R_1) \times T_i$ is an independent set in $\Gamma_0(R)$ for every i : Let $(a_j, b_j) \in \Gamma_0(R_1) \times T_i$ for $j = 1, 2$. If $(a_1, b_1)(a_2, b_2) = (0, 0)$, then $b_1 b_2 = 0$, so that $b_1^2 = 0$ if $b_1 = b_2$ or b_1 is adjacent to b_2 in T_i if $b_1 \neq b_2$. Both statements lead to contradiction.

Notice that by (1), $\Gamma_0(R)$ is a disjoint union of $t_1 + t_2 - 1$ independent sets. Now, we use $t_1 + t_2 - 1$ colors to color those independent sets. In this way, one can easily check that every two adjacent vertices have different colors. Thus, $\chi(\Gamma_0(R)) \leq \chi(\Gamma_0(R_1)) + \chi(\Gamma_0(R_2)) - 1$. $\square$

Similar to Lemma 3.4, we have the following.

Lemma 3.5 *Let R_1 and R_2 be rings and $R \cong R_1 \times R_2$. Then the following hold.*

(i) $\omega(\Gamma_0(R)) \geq \omega(\Gamma_0(R_1)) + \omega(\Gamma_0(R_2)) - 1$.

(ii) If R_2 is reduced, then $\omega(\Gamma_0(R)) = \omega(\Gamma_0(R_1)) + \omega(\Gamma_0(R_2)) - 1$.

Proof. (i) If $\omega(\Gamma_0(R_1))$ or $\omega(\Gamma_0(R_2))$ are infinite then so is $\omega(\Gamma_0(R))$. Therefore the inequality holds trivially. So we may assume that $t_i = \omega(\Gamma_0(R_i)) < \infty$ for $i = 1, 2$. Therefore, there is a maximal clique S (resp. T) in $\Gamma_0(R_1)$ (resp. $\Gamma_0(R_2)$) such that $|S| = t_1$ (resp. $|T| = t_2$). Notice that $\{(0, 0)\} \cup ((S - \{0\}) \times \{0\}) \cup (\{0\} \times (T - \{0\}))$ is a clique in $\Gamma_0(R)$. Thus, $\omega(\Gamma_0(R)) \geq \omega(\Gamma_0(R_1)) + \omega(\Gamma_0(R_2)) - 1$.

(ii) Suppose that R_2 is reduced. As in (i), we may assume that $t_i = \omega(\Gamma_0(R_i)) < \infty$ for $i = 1, 2$. Let W be any maximal clique of $\Gamma_0(R)$. Let $s = |W \cap (\Gamma_0(R_1) \times (\Gamma_0(R_2) - \{0\}))|$ and $\{(a_1, b_1), \dots, (a_s, b_s)\} = W \cap (\Gamma_0(R_1) \times (\Gamma_0(R_2) - \{0\}))$. Notice that $b_i b_j = 0$ for every $i \neq j$. If $b_i = b_j$ for some $i \neq j$, then $b_i^2 = 0$, so that $b_i = 0$ as R_2 is reduced, a contradiction. Therefore $\{0, b_1, \dots, b_s\}$ is a clique of $\Gamma_0(R_2)$. It follows that $s + 1 \leq t_2$. On the other hand, $W \cap (\Gamma_0(R_1) \times \{0\})$ is a clique of $\Gamma_0(R_1) \times \{0\}$. Thus, $|W \cap (\Gamma_0(R_1) \times \{0\})| \leq t_1$. Hence we conclude that $|W| \leq t_1 + t_2 - 1$ and then $\omega(\Gamma_0(R)) \leq \omega(\Gamma_0(R_1)) + \omega(\Gamma_0(R_2)) - 1$. $\square$

An immediately consequence is the following.

Corollary 3.6 *Let $R \cong R_1 \times \dots \times R_k$, where R_i is an integral domain for every i ; then $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = k + 1$.*

We now state and prove our main result in this section.

Theorem 3.7 *Let $R \cong R_1 \times \dots \times R_k$, where R_i is a coloring ring for every i . Suppose that there is a finite set S_i of $\Gamma_0(R_i)$ for every i such that the following hold:*

(i) S_i is a maximal clique of $\Gamma_0(R_i)$.

(ii) If $a, b \in \Gamma_0(R_i) - S_i$, then $a^2 \neq 0$ and $ab \neq 0$.

Let $N_i = \{a \in S_i \mid a^2 = 0\} \subseteq S_i$ and $n_i = |N_i|$ for every i ; then $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = \prod_{i=1}^k n_i + \sum_{i=1}^k (s_i - n_i)$, where $s_i = |S_i|$.

Proof. Rearrange R_i if necessary, we may assume that $S_i = N_i$ if and only if $i > t$ for some nonnegative integer $t \leq k$. For $i = 1, \dots, t$, let

$$A_i = \{(a_1, \dots, a_k) \in \Gamma_0(R) \mid a_j = 0 \text{ if } j \neq i \text{ and } a_i \in S_i - N_i\}$$

and $A = \cup_{i=1}^t A_i$. Let

$$N = N_1 \times \dots \times N_k,$$

$$S = S_1 \times \dots \times S_k,$$

$$T_1 = (Z(R_1) \times \dots \times Z(R_k)) - S$$

and

$$T_2 = (\Gamma_0(R_1) \times \dots \times \Gamma_0(R_k)) - T_1;$$

then $\Gamma_0(R) = S \sqcup T_1 \sqcup T_2$. Further, $N \cup A$ is a clique. Let $n = \prod_{i=1}^k n_i + \sum_{i=1}^k (s_i - n_i)$; then $|N \cup A| = n$ and we require n colors to paint $N \cup A$ as $N \cup A$ is a clique. We will show that $\chi(\Gamma_0(R)) \leq n$. For this, we fix some notations:

For every $i = 1, \dots, k$, choose $b_i \in S_i - \{0\}$. This can be done as $|S_i| \geq 2$.

For every $i = 1, \dots, k$ and $c \in Z(R_i) - S_i$, there are vertices in S_i that are not adjacent to c by assumption (i). Choose an element enjoys this property and denote this element as $\tilde{c}$. So $c\tilde{c} \neq 0$ and $\tilde{c} \in S_i$.

If $(u_1, \dots, u_k) \in N \cup A$ and T is an independent set of $\Gamma_0(R)$, then we use $T \sim \{(u_1, \dots, u_k)\}$ to denote the fact that every vertex of T is colored by the color of $(u_1, \dots, u_k)$. Furthermore, if U and V are two vertices of $\Gamma_0(R)$, then we use $U \sim V$ (resp. $U \approx V$) to denote the fact that the colors of U and V are the same (resp. different).

To finish the proof, we need to color every vertex of S , T_1 and T_2 .

Coloring of S :

First, we use n colors to color $N \cup A$. Notice that

$$S - N = \sqcup_{i=1}^t \sqcup_{a \in S_i - N_i} N_1 \times \dots \times N_{i-1} \times \{a\} \times S_{i+1} \times \dots \times S_k.$$

Notice also that $N_1 \times \cdots \times N_{i-1} \times \{a\} \times S_{i+1} \times \cdots \times S_k$ is an independent set for every $a \in S_i - N_i$ and for every i . Now, we can use the color of $(0, \dots, 0, a, 0 \dots, 0)$ to color the above independent set, that is,

$$N_1 \times \cdots \times N_{i-1} \times \{a\} \times S_{i+1} \times \cdots \times S_k \sim \{(0, \dots, 0, a, 0 \dots, 0)\}. \quad (2)$$

Coloring of T_1 :

Observe that

$$T_1 = \sqcup_{i=1}^k \sqcup_{c \in Z(R_i) - S_i} S_1 \times \cdots \times S_{i-1} \times \{c\} \times Z(R_{i+1}) \times \cdots \times Z(R_k).$$

Moreover, $S_1 \times \cdots \times S_{i-1} \times \{c\} \times Z(R_{i+1}) \times \cdots \times Z(R_k)$ is an independent set for every $c \in Z(R_i) - S_i$ and for every i . Now, we can use the color of $(0, \dots, 0, \tilde{c}, 0 \dots, 0)$ to color the above independent set, that is,

$$S_1 \times \cdots \times S_{i-1} \times \{c\} \times Z(R_{i+1}) \times \cdots \times Z(R_k) \sim \{(0, \dots, 0, \tilde{c}, 0 \dots, 0)\}. \quad (3)$$

Coloring of T_2 :

Observe that

$$T_2 = \sqcup_{i=1}^k \sqcup_{b \in \Gamma_0(R_i) - Z(R_i)} Z(R_1) \times \cdots \times Z(R_{i-1}) \times \{b\} \times \Gamma_0(R_{i+1}) \times \cdots \times \Gamma_0(R_k).$$

Moreover, $Z(R_1) \times \cdots \times Z(R_{i-1}) \times \{b\} \times \Gamma_0(R_{i+1}) \times \cdots \times \Gamma_0(R_k)$ is an independent set for every $b \in \Gamma_0(R_i) - Z(R_i)$ and for every i . Now, we can use the color of $(0, \dots, 0, b_i, 0 \dots, 0)$ to color the above independent set, that is,

$$Z(R_1) \times \cdots \times Z(R_{i-1}) \times \{b\} \times \Gamma_0(R_{i+1}) \times \cdots \times \Gamma_0(R_k) \sim \{(0, \dots, 0, b_i, 0 \dots, 0)\}. \quad (4)$$

To complete the proof, we need to verify that every two adjacent vertices in $\Gamma_0(R)$ have different colors. For this, let $C = (c_1, \dots, c_k)$ and $D = (d_1, \dots, d_k)$ be two adjacent vertices in $\Gamma_0(R)$; then $c_i d_i = 0$ for every i . We proceed by discussing the following cases.

Case 1. $C, D \in S$. Assume first that $C, D \in N$. Then $C \approx D$ of course. Assume next that $C \in N$ and $D \notin N$. Let i be the least integer for which $d_i \notin N_i$; then $D \sim D' = (0, \dots, 0, d_i, 0 \dots, 0)$ by (2). Since $c_i d_i = 0$, it follows that C and D' are different vertices in $N \cup A$. Therefore, $C \approx D'$ and then $C \approx D$. Finally, assume that $C \notin N$ and $D \notin N$. Let i be the least

integer for which $c_i \notin N_i$ and let j be the least integer for which $d_j \notin N_j$; then $C \sim C' = (0, \dots, 0, c_i, 0 \dots, 0)$ and $D \sim D' = (0, \dots, 0, d_j, 0 \dots, 0)$ by (2). If $i = j$, then $c_i d_i = 0$ implies that $C' \approx D'$. Thus, $C \approx D$. If $i \neq j$, then of course C' and D' are different vertices in $N \cup A$, so that $C' \approx D'$, it follows that $C \approx D$.

Case 2. $C \in S$ and $D \in T_1$. Let i be the least integer for which $d_i \notin S_i$; then $D \sim D' = (0, \dots, 0, \tilde{d}_i, 0 \dots, 0)$ by (3). Furthermore, $c_i d_i = 0$ and $d_i \tilde{d}_i \neq 0$ implies that c_i and $\tilde{d}_i$ are different vertices in S_i . Since S_i is a clique, $c_i \tilde{d}_i = 0$. If $C \in N$, then $C \approx D'$. If $C \notin N$, then $C \sim C' = (0, \dots, 0, c_j, 0 \dots, 0)$ by (2), so that C' and D' are different vertices in $N \cup A$, it follows that $C \approx D$. Case 3. $C \in S$ and $D \in T_2$. Let i be the least integer for which $d_i \notin Z(R_i)$; then $D \sim D' = (0, \dots, 0, b_i, 0 \dots, 0)$ by (4). Furthermore, $c_i d_i = 0$ and d_i is a non-zero-divisor implies that $c_i = 0$ and then c_i and b_i are different vertices in S_i . If $C \in N$, then $C \approx D'$. If $C \notin N$, then $C \sim C' = (0, \dots, 0, c_j, 0 \dots, 0)$ for some $j \neq i$ by (2), so that C' and D' are different vertices in $N \cup A$, it follows that $C \approx D$.

Case 4. $C, D \in T_1$. In this case, let i be the least integer for which $c_i \notin S_i$ and let j be the least integer for which $d_j \notin S_j$; then $C \sim C' = (0, \dots, 0, \tilde{c}_i, 0 \dots, 0)$ and $D \sim D' = (0, \dots, 0, \tilde{d}_j, \dots, 0)$ by (3). If $i = j$, then $c_i d_i = 0$, contradicts to the assumption (ii). Therefore, $i \neq j$. It follows that C' and D' are different vertices in $N \cup A$, $C' \approx D'$. Thus, $C \approx D$.

Case 5. $C \in T_1$ and $D \in T_2$. In this case, let i be the least integer for which $c_i \notin S_i$ and let j be the least integer for which $d_j \notin Z(R_j)$; then $C \sim C' = (0, \dots, 0, \tilde{c}_i, 0 \dots, 0)$ by (3) and $D \sim D' = (0, \dots, 0, b_j, \dots, 0)$ by (4). If $i = j$, then $c_i d_i = 0$, contradicts to the assumption (ii). Therefore, $i \neq j$. It follows that C' and D' are different vertices in $N \cup A$, $C' \approx D'$. Thus, $C \approx D$.

Case 6. $C, D \in T_2$. In this case, let i be the least integer for which $c_i \notin Z(R_i)$ and let j be the least integer for which $d_j \notin Z(R_j)$; then $C \sim C' = (0, \dots, 0, b_i, 0 \dots, 0)$ and $D \sim D' = (0, \dots, 0, b_j, \dots, 0)$ by (4). If $i = j$, then $c_i d_i = 0$, contradicts to the assumption (ii). Therefore, $i \neq j$. It follows that C' and D' are different vertices in $N \cup A$, $C' \approx D'$. Thus, $C \approx D$. $\square$

Corollary 3.8 *Let $R \cong \mathbb{Z}^n \times \mathbb{Z}_{p_1^{r_1}} \times \dots \times \mathbb{Z}_{p_k^{r_k}}$; then $\chi(\Gamma_0(R)) = \omega(\Gamma_0(R)) = \prod_{i=1}^k p_i^{\lfloor r_i/2 \rfloor} + t + n$, where $\lfloor n \rfloor$ is the largest integer that is smaller than or equal*

to n and $t = |\{i \mid r_i \text{ is odd}\}|$.

Proof. By Lemma 3.4, Lemma 3.5 and Corollary 3.6, we may assume that $R \cong \mathbb{Z}_{p_1^{r_1}} \times \cdots \times \mathbb{Z}_{p_k^{r_k}}$. Let $S_i = \{p_i^{r_i/2}n \mid n \in \mathbb{Z}\}$ if r_i is even and $S_i = \{p_i^{(r_i+1)/2}n \mid n \in \mathbb{Z}\} \cup \{p_i^{(r_i-1)/2}\}$ if r_i is odd; then S_i satisfies (i) and (ii) of Theorem 3.7. Let $N_i = \{a \in S_i \mid a^2 = 0\}$; then by Lemma 3.1 and Corollary 3.6, $|S_i - N_i| = 1$ and $|N_i| = p^{\lfloor r_i/2 \rfloor}$ if r_i is odd. Further, $|S_i - N_i| = 0$ and $|N_i| = p^{\lfloor r_i/2 \rfloor}$ if r_i is even. Thus, the assertion follows. $\square$

Since $\Gamma(R)$ is a subgraph of $\Gamma_0(R)$ and $\Gamma_0(R) - \{0\}$ is a disjoint union of $\Gamma(R)$ and a totally disconnected graph. We have the following.

Corollary 3.9 *Let $R \cong \mathbb{Z}_{p_1^{r_1}} \times \cdots \times \mathbb{Z}_{p_k^{r_k}}$; then $\chi(\Gamma(R)) = \omega(\Gamma(R))$.*

Remark 3.10 (i) In [5, Theorem 3.2], Anderson and Naseer also provide some sufficient conditions for the product of coloring rings to be chromatic. However, [5, Theorem 3.2] can not derive Theorem 3.7. For example, if $R \cong \mathbb{Z}^n$, then Theorem 3.7 can be used to obtain that R is chromatic. However, [5, Theorem 3.2] can not.

(ii) Theorem 3.7 can be used to derive [5, Corollary 3.3].

(iii) In [6], it is shown that if $R \cong \mathbb{Z}_n$, then R is chromatic. Corollary 3.8 generalize this result.

4 Coloring of $\overline{\Gamma(R)}$

Throughout, let $R \cong \mathbb{Z}_{p_1^{r_1}} \times \cdots \times \mathbb{Z}_{p_k^{r_k}}$. Let $\Gamma(R)$ be the graph associated to R defined in the introduction. For convenience, we fix some notion from [8].

Definition 4.1 *Let $R \cong \mathbb{Z}_{p^r}$, where p is a prime number and r be a positive integer. Let a, b be two distinct vertices of $\Gamma(R)$; then a and b are associated, denoted by $a \sim b$, if there is a unit u in R such that $a = ub$. The associate class of a , denoted by A_a , is the set $\{b \in \Gamma(R) \mid b \sim a\}$. Thus, $\Gamma(R) = \sqcup_{i=1}^r A_{p^i}$, $1 = |A_{p^r}| < |A_{p^{r-1}}| < \cdots < |A_{p^1}|$ and $|A_{p^i}| = p^{r-i} - p^{r-i-1}$ for $i = 0, 1, \dots, r-1$.*

Lemma 4.2 *Let $R \cong \mathbb{Z}_{p^r}$, where p is a prime number and $r \geq 2$ be a positive integer. Then $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)}) = p^{r-1} - p^{r-(r/2)}$ if r is even and $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)}) = p^{r-1} - p^{(r-1)/2}$ if r is odd.*

Proof. If $r = 2$, then $\overline{\Gamma(R)}$ is discrete, so that $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)}) = 0$, therefore the assertion holds. Thus, we may assume that $r \geq 3$. Suppose that r is even. Let $C = \sqcup_{i=1}^{r/2-1} A_{p^i}$; then $|C| = p^{r-1} - p^{r-(r/2)}$ by Definition 4.1 and C is a clique of $\overline{\Gamma(R)}$. Let $n = |C|$; then we require n colors to paint C . Since $|\overline{\Gamma(R)} - C| = p^{r-(r/2)} - 1 \leq n$, we can use part of the colors of C to color $\overline{\Gamma(R)} - C$. Hence $\chi(\overline{\Gamma(R)}) \leq |C| \leq \omega(\overline{\Gamma(R)})$. Suppose that r is odd. Let $C = \sqcup_{i=1}^{(r-1)/2} A_{p^i}$; then $|C| = p^{r-1} - p^{(r-1)/2}$ by Definition 4.1 and C is a clique of $\overline{\Gamma(R)}$. Let $n = |C|$; then we require n colors to paint C . Since $|\overline{\Gamma(R)} - C| = p^{(r-1)/2} - 1 \leq n$, we can use part of the colors of C to color $\overline{\Gamma(R)} - C$. Hence $\chi(\overline{\Gamma(R)}) \leq |C| \leq \omega(\overline{\Gamma(R)})$. $\square$

Example 4.3 Let $R \cong \mathbb{Z}_8 \times \mathbb{Z}_{16}$; then $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)}) = 76$.

Proof. Let

$$C = \{0, 2, 4, 6\} \times \{1, 3, 5, 7, 9, 11, 13, 15\} \\ \sqcup (\mathbb{Z}_8 \times \{2, 6, 10, 14\}) \\ \sqcup (\{1, 3, 5, 7, 2, 6\} \times \{4, 12\})$$

then C is a clique in $\overline{\Gamma(R)}$ as C is an independent set in $\Gamma(R)$. Moreover, C is maximal. Observe that $|\Gamma(R)| = 8 \cdot 16 - 4 \cdot 8 - 1 = 95$ and $|C| = 76$. So, $|\overline{\Gamma(R)} - C| = 19 < |C|$. Now, we require 76 colors to paint C and then we can use part of the colors of C to color $\overline{\Gamma(R)} - C$. Hence, $\chi(\overline{\Gamma(R)}) \leq |C| \leq \omega(\overline{\Gamma(R)})$. $\square$

Theorem 4.4 Let $R \cong \mathbb{Z}_{p_1^{r_1}} \times \cdots \times \mathbb{Z}_{p_k^{r_k}}$, where p_i is a prime number for every i . Then $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)})$.

Proof. By Lemma 4.2, we may assume that $k \geq 2$. Let

$$T = \{(t_1, \dots, t_k) \in \mathbb{N}^k \mid 0 \leq t_i \leq r_i \ \forall i\} - \{(0, \dots, 0), (r_1, \dots, r_k)\}.$$

For any k -tuple $(t_1, \dots, t_k) \in T$, let

$$A(t_1, \dots, t_k) = A_{p_1^{t_1}} \times \cdots \times A_{p_k^{t_k}}.$$

Here, $A_{p_i^{t_i}}$ is the associated class of $p_i^{t_i}$ in $\mathbb{Z}_{p_i^{r_i}}$. It follows that

$$|A(t_1, \dots, t_k)| = \prod_{i=1}^k |A_{p_i^{t_i}}|$$

and

$$\Gamma(R) = \sqcup_{(t_1, \dots, t_k) \in T} A(t_1, \dots, t_k).$$

For any k -tuple $(t_1, \dots, t_k) \in T$, define

$$\tilde{A}(t_1, \dots, t_k) = A_{p_1^{r_1-t_1}} \times \dots \times A_{p_k^{r_k-t_k}} = A(r_1-t_1, \dots, r_k-t_k).$$

Observe that if r_i is even for every i , then $\tilde{A}(r_1/2, \dots, r_k/2) = A(r_1/2, \dots, r_k/2)$. If this is the case, we use A for $A(r_1/2, \dots, r_k/2)$.

Consider the two following subsets of T :

$$T_1 = \{(t_1, \dots, t_k) \in T \mid |A(t_1, \dots, t_k)| > |\tilde{A}(t_1, \dots, t_k)|\}$$

and

$$T_2 = \{(t_1, \dots, t_k) \in T \mid |A(t_1, \dots, t_k)| = |\tilde{A}(t_1, \dots, t_k)| \text{ and there is an } i \text{ such that } t_i \neq r_i/2\}$$

Let T_0 be the subset of T consists of every k -tuple of T_1 and exactly one k -tuple of $\{(t_1, \dots, t_k), (r_1-t_1, \dots, r_k-t_k)\}$ from T_2 . Notice that $(r_1/2, \dots, r_k/2) \notin T_0$ if r_i is even for every i . Notice also that if $(t_1, \dots, t_k) \in T_0$, then there is an integer i such that $t_i < r_i/2$. For if not, then $t_i \geq r_i/2$ for every i . Therefore $|A_{p_i^{t_i}}| \leq |A_{p_i^{r_i-t_i}}|$ for every i . If $t_j \neq r_j/2$, then $t_j > r_j/2$ for some j , so that $|A_{p_j^{t_j}}| < |A_{p_j^{r_j-t_j}}|$, it follows that $|A(t_1, \dots, t_k)| < |\tilde{A}(t_1, \dots, t_k)|$, a contradiction. Thus, r_i is even and $t_i = r_i/2$ for every i , contradiction again.

Let V be a vertex in A if r_i is even for every i . Further, let

$$C = \sqcup_{(t_1, \dots, t_k) \in T_0} A(t_1, \dots, t_k) \sqcup \{V\}$$

if r_i is even for every i and

$$C = \sqcup_{(t_1, \dots, t_k) \in T_0} A(t_1, \dots, t_k)$$

if r_i is odd for some i .

In the following, we will show that C is a clique in $\overline{\Gamma(R)}$. For this, we first examine that $A(t_1, \dots, t_k)$ is a clique in $\overline{\Gamma(R)}$ if $(t_1, \dots, t_k) \in T_0$. Or equivalently, to show that $A(t_1, \dots, t_k)$ is an independent set in $\Gamma(R)$. Suppose not. Then there are vertices $(a_1, \dots, a_k), (b_1, \dots, b_k) \in A(t_1, \dots, t_k)$ such that $a_i b_i = 0$ for every i . However, this implies that $t_i \geq r_i/2$ for every i , contradicts to the fact that there is an integer i such that $t_i < r_i/2$ if $(t_1, \dots, t_k) \in T_0$.

We next show that if $(c_1, \dots, c_k) = C \in A(t_1, \dots, t_k)$ and $(d_1, \dots, d_k) = D \in A(s_1, \dots, s_k)$, then $(c_1, \dots, c_k)$ and $(d_1, \dots, d_k)$ are adjacent in $\overline{\Gamma(R)}$, where $(t_1, \dots, t_k)$ and $(s_1, \dots, s_k)$ are different k -tuples in T_0 . Or equivalently, to show that $(c_1, \dots, c_k)$ is not adjacent to $(d_1, \dots, d_k)$ in $\Gamma(R)$. Suppose not. Then $c_i d_i = 0$ for every i . This implies that $t_i + s_i \geq r_i$ for every i . Since $A(s_1, \dots, s_k) \neq \tilde{A}(t_1, \dots, t_k)$, the set $\{i \mid t_i + s_i > r_i\}$ is nonempty. We may assume that $t_1 + s_1 > r_1$. Therefore, $t_1 \geq 1$ and $s_1 \geq 1$. For, if $t_1 = 0$ or $s_1 = 0$, then c_1 or d_1 is a unit, so that $s_1 = r_1$ or $t_1 = r_1$, it follows that $t_1 + s_1 = r_1$, a contradiction. Thus,

$$|A(t_1, t_2, \dots, t_k)| < |A(t_1-1, t_2, \dots, t_k)| \leq |A(r_1-s_1, \dots, r_k-s_k)| = |\tilde{A}(s_1, \dots, s_k)|$$

and

$$|A(s_1, s_2, \dots, s_k)| < |A(s_1-1, s_2, \dots, s_k)| \leq |A(r_1-t_1, \dots, r_k-t_k)| = |\tilde{A}(t_1, \dots, t_k)|.$$

Thus, $|A(t_1, \dots, t_k)| < |A(s_1, \dots, s_k)|$ and $|A(s_1, \dots, s_k)| < |A(t_1, \dots, t_k)|$, a contradiction.

Now, if r_i is even for every i , then A is a clique in $\Gamma(R)$, so that V is not adjacent to every vertex of $A - \{V\}$ and $A - \{V\}$ is an independent set in $\overline{\Gamma(R)}$. Moreover, V is adjacent to every vertex of $A(t_1, \dots, t_k)$ in $\overline{\Gamma(R)}$ if $(t_1, \dots, t_k) \in T_0$ as there is an integer i such that $t_i < r_i/2$ and then V is not adjacent to every vertex of $A(t_1, \dots, t_k)$ in $\Gamma(R)$.

From the above, we see that C is a clique in $\overline{\Gamma(R)}$. Let $n = |C|$; then we require n different colors to paint C . We then use the colors of $A(t_1, \dots, t_k)$ ($(t_1, \dots, t_k) \in T_0$) to color $\tilde{A}(t_1, \dots, t_k)$ and the color of V to color every vertex of $A - \{V\}$ if r_i is even for every i . In this way, it is easy to see that every two adjacent vertices in $\overline{\Gamma(R)}$ have different colors. Hence, we conclude that $\chi(\overline{\Gamma(R)}) \leq n \leq \omega(\overline{\Gamma(R)})$. $\square$

An easy consequence of Theorem 4.4 is that if $R \cong \mathbb{Z}_{p_1} \times \dots \times \mathbb{Z}_{p_k}$, then $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)})$. However, from the proof of Theorem 4.4, one can replace each $\mathbb{Z}_{p_i}$ by any finite field.

Corollary 4.5 *Let $R \cong F_1 \times \dots \times F_k$, where F_i is a finite fields for every i . Then $\chi(\overline{\Gamma(R)}) = \omega(\overline{\Gamma(R)})$.*

References

- [1] S. Akbari, H.R. Maimani and S. Yassemi, *When a zero-divisor graph is planar or a complete r -partite graph*, J. Algebra **270** (2003), 169–180.

- [2] S. Akbari and A. Mohammadian, *On the zero-divisor graph of a commutative ring*, J. Algebra **274** (2004), 847–855.
- [3] D.F. Anderson, A. Frazier, A. Lauve and P.S. Livingston, *The zero-divisor graph of a commutative ring, II*, Lecture Notes in Pure and Appl. Math. **220** (2001), 61–72.
- [4] D.F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434–447.
- [5] D.D. Anderson and M. Naseer, *Beck's coloring of a commutative ring*, J. Algebra **159** (1993), 500–514.
- [6] I. Beck, *Coloring of commutative rings*, J. Algebra **116** (1988), 208–226.
- [7] M. Brickel, *Complements of zero-divisor graphs*, Plan B Project, Miami university, 2004.
- [8] N.I. Córdova, C. Gholston and H.A. Hauser, *The structure of zero-divisor graphs*, preprint.
- [9] N.O. Smith, *Planar zero-divisor graphs*, International J. Commutative Rings **2** (2003), no. 4, 177–188.
- [10] H.-J. Wang, *Zero-divisor graphs of genus one*, J. Algebra **304** (2006), no. 2, 666–678.

Department of Mathematics,
National Chung Cheng University,
Chiayi 621, Taiwan
hjjwang@math.ccu.edu.tw